



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

**Cultivation and Genomic Analysis of “Candidatus Nitrosocaldus islandicus,” an Obligately Thermophilic, Ammonia-Oxidizing Thaumarchaeon from a Hot Spring Biofilm in Graendalur Valley, Iceland**

Daebeler, Anne; Herbold, Craig; Vierheilig, Julia; J. Sedlacek, Christopher; Pjevac, Petra; Albertsen, Mads; Kirkegaard, Rasmus Hansen; R. de la Torre, Jose; Daims, Holger; Wagner, Michael

*Published in:*  
Frontiers in Microbiology

*DOI (link to publication from Publisher):*  
[10.3389/fmicb.2018.00193](https://doi.org/10.3389/fmicb.2018.00193)

*Creative Commons License*  
CC BY 4.0

*Publication date:*  
2018

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Daebeler, A., Herbold, C., Vierheilig, J., J. Sedlacek, C., Pjevac, P., Albertsen, M., Kirkegaard, R. H., R. de la Torre, J., Daims, H., & Wagner, M. (2018). Cultivation and Genomic Analysis of “Candidatus Nitrosocaldus islandicus,” an Obligately Thermophilic, Ammonia-Oxidizing Thaumarchaeon from a Hot Spring Biofilm in Graendalur Valley, Iceland. *Frontiers in Microbiology*, 9(FEB), [193]. <https://doi.org/10.3389/fmicb.2018.00193>

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -



# Cultivation and Genomic Analysis of “*Candidatus Nitrosocaldus islandicus*,” an Obligately Thermophilic, Ammonia-Oxidizing Thaumarchaeon from a Hot Spring Biofilm in Graendalur Valley, Iceland

## OPEN ACCESS

### Edited by:

Peter Dunfield,  
University of Calgary, Canada

### Reviewed by:

Jeremy Dodsworth,  
California State University,  
San Bernardino, United States  
Nathan Basiliko,  
Laurentian University, Canada  
Martin G. Klotz,  
Washington State University Tri-Cities,  
United States

### \*Correspondence:

Anne Daebeler  
daebeler@microbial-ecology.net  
Michael Wagner  
wagner@microbial-ecology.net

### Specialty section:

This article was submitted to  
Extreme Microbiology,  
a section of the journal  
Frontiers in Microbiology

**Received:** 15 December 2017

**Accepted:** 29 January 2018

**Published:** 14 February 2018

### Citation:

Daebeler A, Herbold CW, Vierheilig J, Sedlacek CJ, Pjevac P, Albertsen M, Kirkegaard RH, de la Torre JR, Daims H and Wagner M (2018) Cultivation and Genomic Analysis of “*Candidatus Nitrosocaldus islandicus*,” an Obligately Thermophilic, Ammonia-Oxidizing Thaumarchaeon from a Hot Spring Biofilm in Graendalur Valley, Iceland. *Front. Microbiol.* 9:193. doi: 10.3389/fmicb.2018.00193

Anne Daebeler<sup>1\*</sup>, Craig W. Herbold<sup>1</sup>, Julia Vierheilig<sup>1</sup>, Christopher J. Sedlacek<sup>1</sup>, Petra Pjevac<sup>1</sup>, Mads Albertsen<sup>2</sup>, Rasmus H. Kirkegaard<sup>2</sup>, José R. de la Torre<sup>3</sup>, Holger Daims<sup>1</sup> and Michael Wagner<sup>1\*</sup>

<sup>1</sup> Division of Microbial Ecology, Department of Microbiology and Ecosystem Science, Research Network “Chemistry meets Microbiology”, University of Vienna, Vienna, Austria, <sup>2</sup> Department of Chemistry and Bioscience, Center for Microbial Communities, Aalborg University, Aalborg, Denmark, <sup>3</sup> Department of Biology, San Francisco State University, San Francisco, CA, United States

Ammonia-oxidizing archaea (AOA) within the phylum Thaumarchaeota are the only known aerobic ammonia oxidizers in geothermal environments. Although molecular data indicate the presence of phylogenetically diverse AOA from the *Nitrosocaldus* clade, group 1.1b and group 1.1a Thaumarchaeota in terrestrial high-temperature habitats, only one<sup>§</sup> enrichment culture of an AOA thriving above 50°C has been reported and functionally analyzed. In this study, we physiologically and genomically characterized a newly discovered thaumarchaeon from the deep-branching Nitrosocaldaceae family of which we have obtained a high (~85%) enrichment from biofilm of an Icelandic hot spring (73°C). This AOA, which we provisionally refer to as “*Candidatus Nitrosocaldus islandicus*,” is an obligately thermophilic, aerobic chemolithoautotrophic ammonia oxidizer, which stoichiometrically converts ammonia to nitrite at temperatures between 50 and 70°C. “*Ca. N. islandicus*” encodes the expected repertoire of enzymes proposed to be required for archaeal ammonia oxidation, but unexpectedly lacks a *nirK* gene and also possesses no identifiable other enzyme for nitric oxide (NO) generation<sup>§</sup>. Nevertheless, ammonia oxidation by this AOA appears to be NO-dependent as “*Ca. N. islandicus*” is, like all other tested AOA, inhibited by the addition of an NO scavenger. Furthermore, comparative genomics revealed that “*Ca. N. islandicus*” has the potential for aromatic amino acid fermentation as its genome encodes an indolepyruvate oxidoreductase (*iorAB*) as well as a type 3b hydrogenase, which are not present in any other sequenced AOA. A further surprising genomic feature of this thermophilic ammonia oxidizer is the absence of DNA polymerase D genes<sup>§</sup> – one of the predominant replicative DNA polymerases in all

<sup>§</sup> See section “Note Added in Proof”.

other ammonia-oxidizing Thaumarchaeota. Collectively, our findings suggest that metabolic versatility and DNA replication might differ substantially between obligately thermophilic and other AOA.

**Keywords:** AOA, thaumarchaeota, thermophile, nitrification, hot spring, *nirK*, polymerase, anaerobic metabolism

## INTRODUCTION

Thaumarchaeota (Brochier-Armanet et al., 2008) are among the most abundant archaeal organisms on Earth, and thrive in most oxic environments (Francis et al., 2007; Erguder et al., 2009; Schleper and Nicol, 2010; Bouskill et al., 2012; Prosser and Nicol, 2012; Stahl and de la Torre, 2012; Stieglmeier et al., 2014a), but have also been detected in anoxic systems (Molina et al., 2010; Bouskill et al., 2012; Buckles et al., 2013; Beam et al., 2014; Lin et al., 2015). This phylum comprises ammonia-oxidizing archaea (AOA) and other archaeal taxa in which ammonia oxidation has not been demonstrated. All cultured members of the Thaumarchaeota are AOA and grow by using ammonia, urea or cyanate as substrate (Palatinszky et al., 2015; Bayer et al., 2016; Sauder et al., 2017; Qin et al., 2017a), although *in situ* experiments suggest that certain members of this phylum capable of ammonia oxidation also possess other lifestyles (Mussmann et al., 2011; Sauder et al., 2017). In aquatic and terrestrial environments Thaumarchaeota often co-occur with ammonia-oxidizing bacteria (AOB), and frequently outnumber them by orders of magnitude (Francis et al., 2005; Leininger et al., 2006; Mincer et al., 2007; Adair and Schwartz, 2008; Abell et al., 2010; Mussmann et al., 2011; Zeglin et al., 2011; Daebeler et al., 2012). Thaumarchaeota also inhabit extreme environments like terrestrial hot springs and other high temperature habitats, where AOB are not detectable (Weidler et al., 2007; Reigstad et al., 2008; Wang et al., 2009; Zhao et al., 2011; Chen et al., 2016). In addition to the presence of Thaumarchaeota in hot environments, high *in situ* nitrification rates (Reigstad et al., 2008; Dodsworth et al., 2011; Chen et al., 2016) and transcription of genes involved in archaeal ammonia oxidation in several hot springs over 74°C (Zhang et al., 2008; Jiang et al., 2010) support an important role of thermophilic AOA in these systems.

Despite their apparent importance for nitrogen cycling in a wide range of thermal habitats, only one<sup>§</sup> thermophilic AOA species from an enrichment culture has been reported to date (de la Torre et al., 2008; Qin et al., 2017b) and was named “*Candidatus* (*Ca.*) *Nitrosocaldus yellowstonensis*.” The authors note that throughout this study the definition for thermophiles by Stetter (1998) as organisms that grow optimally above 50°C is used. In addition, several enrichment cultures and one pure culture of moderately thermophilic AOA, which are able to grow at 50°C, but have growth optima only at temperatures below 50°C, have been described (Hatzenpichler et al., 2008; Lebedeva et al., 2013; Palatinszky et al., 2015). Therefore, our current knowledge on specific adaptations or metabolic capabilities of thermophilic AOA growing preferably at temperatures above 50°C is very limited (Spang et al., 2012).

In 16S rRNA and ammonia monooxygenase subunit A (*amoA*) gene trees “*Ca. Nitrosocaldus yellowstonensis*” branches most deeply among Thaumarchaeota that possess ammonia monooxygenase (AMO) genes. In consequence, the Nitrosocaldales clade has been considered as being close to the evolutionary origin of Thaumarchaeota encoding the genetic repertoire for ammonia oxidation (de la Torre et al., 2008; Spang et al., 2017). However, since the genome sequence of “*Ca. N. yellowstonensis*” is not yet published, phylogenomic analysis to confirm an ancestral position of the Nitrosocaldales relative to other Thaumarchaeota have been pending.

Here we report on the enrichment, phylogenomic analyses, and selected (putative) metabolic features of an obligately thermophilic AOA from the Nitrosocaldales clade obtained from a biofilm collected from a 73°C hot spring in Iceland. This organism, provisionally referred to as “*Ca. Nitrosocaldus islandicus*,” occupies a fundamentally different niche compared to other genomically characterized AOA as its ammonia-oxidizing activity is restricted to temperatures ranging from 50 to 70°C.

## MATERIALS AND METHODS

### Enrichment, Cultivation, and Physiological Experiments

The enrichment of “*Ca. N. islandicus*” was initiated by inoculation of 40 ml sterile mineral medium (Koch et al., 2015) containing 0.5 mM filter-sterilized NH<sub>4</sub>Cl with approximately 0.1 g of hot spring biofilm, which had been submerged in running water at the sampling site in a geothermal area in Graendalur valley, (64° 1' 7" N, 21° 11' 20" W) Iceland. At the sampling site, the spring had a pH of 6.5 and a temperature of 73°C. The culture was initially incubated without agitation in 100 ml glass bottles in the dark at 60°C and checked weekly for ammonium and nitrite content of the medium by using Nessler's reagent (K<sub>2</sub>HgI<sub>4</sub> – KOH solution; Sigma–Aldrich) and nitrite/nitrate test stripes (Merckoquant; Merck). Ammonium (1 mM NH<sub>4</sub>Cl) was replenished when completely consumed. At the same time pH was monitored by using pH test stripes (Macherey–Nagel) and kept at pH 7–8 by titration with NaHCO<sub>3</sub>. When the pH dropped below 6 the enrichment culture ceased to oxidize ammonia, but activity was restored by readjusting the pH to between 7 and 8. The ammonium and nitrite concentrations were quantified photometrically (Kandeler and Gerber, 1988; Miranda et al., 2001) using an Infinite 200 Pro spectrophotometer (Tecan Group AG). The microbial community composition of the enrichment was regularly monitored by fluorescence *in situ* hybridization (FISH) with 16S rRNA-targeted probes labeled with dyes Cy3, Cy5, or Fluos as described elsewhere (Daims et al., 2005). Probes targeting most bacteria (EUB338 probe mix;

<sup>§</sup> See section “Note Added in Proof”.

Amann et al., 1990; Daims et al., 1999), most archaea (Arch915, Stahl and Amann, 1991) and most Thaumarchaeota (Thaum726, Beam, 2015) were applied. All positive results were verified using the nonsense probe non-EUB338 (Wallner et al., 1993) labeled with the same dyes. Additions of the antibiotics rifampicine, gentamicine, and kanamycine at concentrations ranging from 10 to 100 mg l<sup>-1</sup> were applied alone and in combination, but no ammonia oxidizing activity was observed in the course of 8 weeks. Treatments with the macrolide antibiotic spiramycin (15 mg l<sup>-1</sup>), which partly retains its antibacterial activity at 60°C (Zorraquino et al., 2011), were performed as described in Zhang et al. (2015) together with serial dilutions ranging from 10<sup>-5</sup> to 10<sup>-8</sup> to obtain a highly enriched (~85%) AOA culture that was used for further characterization.

Growth rates of “*Ca. N. islandicus*” were determined across a range of incubation temperatures (50–70°C). Triplicate cultures (25 ml) and negative controls (cultures not supplied with ammonium or inoculated with autoclaved biomass) were incubated for 10 days in 100 ml Schott bottles without agitation in the dark at the respective temperature. Samples from these experiments were either stored at –20°C for subsequent qPCR analyses (150 µl) or centrifuged (21,000 × g, 15 min, 18°C) to remove cells and the supernatant was stored at –20°C for chemical analysis (600 µl). qPCR analysis with primers CrenamoA19F (Leininger et al., 2006) and CrenamoA616R (Tourna et al., 2008) targeting the archaeal *amoA* gene was otherwise performed as described in Pjevac et al. (2017) before the genome sequence of “*Ca. N. islandicus*” was available. However, subsequent analysis demonstrated that the employed qPCR primers contain mismatches to the *amoA* sequence of this AOA in the middle of the forward and reverse primer. Average cell numbers were calculated from samples of triplicate cultures ( $n = 3 \pm$  standard error). The specific growth rate was calculated from log-linear plots of *amoA* gene abundance in individual cultures. In this analysis, three out of seven time points were interpolated through linear regression.

To test whether the NO-scavenger 2-phenyl-4,4,5,5-tetramethylimidazoline-3-oxide-1-oxyl (PTIO; TCI, Germany) inhibits ammonia oxidation by “*Ca. N. islandicus*,” 40 ml aliquots of mineral medium containing 1 mM ammonium were inoculated with 10% (v/v) of an exponential-phase culture and incubated in duplicates in the presence of 0, 33, and 100 µM PTIO, respectively. PTIO was dissolved in sterile mineral medium before addition to the cultures. The cultures not exposed to PTIO were supplemented with the same volume of sterile medium. The cultures were sampled (2 ml) at the beginning of the experiment and after 15 days of incubation. Nitrite and ammonium concentrations were measured as described above.

## DNA Extraction, Genome Sequencing, and Annotation

DNA from three replicate enrichment cultures containing “*Ca. N. islandicus*” as the only detectable ammonia oxidizer was extracted as described by Angel and Conrad (2013) and sequenced by Illumina HiSeq next generation sequencing (250 bp paired end reads). Since we did not obtain a complete genome with this

approach we extracted genomic DNA from the enrichment at a later stage according to Zhou et al. (1996) yielding high molecular weight DNA. Genomic DNA was then sheared in a Covaris g-TUBE (Covaris, United States) at 9000 RPM for 2x 1 min. in an Eppendorf mini spin plus centrifuge (Eppendorf, DE). The DNA was run on a E-Gel EX 1% agarose gel (Thermo Fisher, United States) and small DNA fragments were removed by excising a band with a length of ~8 kb. The DNA was purified from the gel cut using the ultraClean 15 DNA Purification Kit (Qiagen, United States). The DNA was prepared for sequencing using the “1D Low Input gDNA with PCR SQK-LSK108” protocol (Oxford Nanopore Technologies, United Kingdom) and sequenced on a FLO-MIN106 flowcell using the MinION MK1b (Oxford Nanopore Technologies, United Kingdom) following the manufacturers protocol using MinKNOW (v. 1.7.14). The nanopore reads were basecalled using Albacore (V. 2.0.1) (Oxford Nanopore Technologies, United Kingdom). The taxonomy was assigned to contigs as described in Karst et al. (unpublished). The read coverage was calculated from the read mappings. The complete genome was assembled using a hybrid approach combining the data from the Illumina and Nanopore sequencing with the hybrid assembler Unicycler (v. 0.4.1, Wick et al., 2017). The genome bins of the two contaminating organisms were assembled from the Nanopore reads using Miniasm (Li, 2016) and polished twice with the Nanopore reads using Racon (Vaser et al., 2017). No other microbe, encoding genes indicative for ammonia-oxidation, was identified in either of the two metagenomes.

The complete genome of “*Ca. N. islandicus*” was uploaded to the MicroScope platform (Vallenet et al., 2013) for automatic annotation, which was amended manually where necessary. The full genome sequence of “*Ca. N. islandicus*” has been deposited in GenBank (accession CP024014) and associated annotations are publicly available in MicroScope (“*Candidatus Nitrosocaldus islandicus* strain 3F”). The raw sequence (Illumina MiSeq and Nanopore) data has been deposited to ENA under Study accession number PRJEB24462.

Protein-coding genes from the thaumarchaeon studied here were compared to those from 30 Thaumarchaeota with available genomic data (Supplementary Table S1) downloaded from NCBI. The coding sequences (CDS) with accession numbers from each genome, as downloaded from NCBI, were combined with additional CDS predictions made by Prodigal (Hyatt et al., 2010) to account for variability in CDS predictions from different primary data providers and platforms. Predicted CDS from the thaumarchaeon of this study were aligned to CDS from reference genomes using blastp (Word\_size = 2, substitution matrix BLOSUM45). Genes were considered homologous only if the blastp alignment exceeded 50% of the length of both query and subject sequences. CDS of “*Ca. N. islandicus*” that lacked any homologs in other Thaumarchaeota were considered “unique.” Unique CDS of unknown function were searched for secretion signals and for predicted membrane-spanning domains of the encoded proteins using the Phobius web server (Käll et al., 2007) and putative structures were determined using the Phyre2 web server (Kelley et al., 2015). Homology to “Thaumarchaeota-core” proteins was assessed by cross-referencing the blastp homology



search to the proteins defined for “*Ca. Nitrosotalea devanaterre*” by Herbold et al. (2017).

## Phylogenetic Analysis and Habitat Preference

For 16S rRNA and *amoA* gene-based phylogenetic analysis, the full-length 16S rRNA and *amoA* gene sequences of “*Ca. N. islandicus*” retrieved from the genome assembly were imported into the ARB software package (Ludwig et al., 2004) together with other full length 16S rRNA or *amoA* gene sequences from cultivated AOA strains and aligned with the integrated ARB aligner with manual curation. 171 sequences from the Aigarchaea were included in the alignment and used as outgroup in the 16S rRNA gene phylogenetic analyses. For the *amoA* gene phylogenetic analyses no outgroup was selected. The 16S rRNA and *amoA* gene consensus trees were reconstructed using Maximum-Likelihood (ML; using the GTRGAMMA evolution model), Neighbour Joining (NJ) and Maximum Parsimony (MP) methods. For all calculations, a sequence filter considering only positions conserved in  $\geq 50\%$  of all thaumarchaeotal and aigarchaeal sequences was used, resulting in 2444 and 488 alignment positions for the 16S rRNA and *amoA* genes, respectively.

A Bayesian-inference phylogenomic tree was obtained using the automatically generated alignment of 34 concatenated universal marker genes (Supplementary Table S2), which were identified by CheckM in Parks et al. (2015). This alignment was used as input for PhyloBayes (Lartillot et al., 2009) with 10 independent chains of 4,000 generations using the CAT-GTR model; 2,000 generations of each chain were discarded as burn-in, the remainder were subsampled every second tree (bpcomp -x 2000 2 4000) and pooled together for calculation of posterior probabilities.

Whole-genome based average nucleotide identity (gANI, Varghese et al., 2015) and average amino acid identity values (AAI, Konstantinidis and Tiedje, 2005) were calculated between the genomes of “*Ca. N. islandicus*” and “*Ca. N. yellowstonensis*” using sets of annotated genes supplemented with additional gene calls predicted by Prodigal (Hyatt et al., 2010). gANI was calculated using the Microbial Species Identifier (MiSI) method (Varghese et al., 2015). For AAI, bidirectional best hits were identified using blastp, requiring that query genes aligned over at least 70% of their length to target genes (in each unidirectional blastp search). Query gene length was used to calculate a weighted average % identity over all best hit pairs and the calculations were repeated using each genome as query and target.

The occurrence of organisms closely related to “*Ca. N. islandicus*” and “*Ca. N. yellowstonensis*” in publicly deposited amplicon sequencing data sets was assessed using IMNGS (Lagkouvardos et al., 2016) with the full-length 16S rRNA gene sequences of both organisms as query and a nucleotide identity threshold of 97%.

*PolB* amino acid sequences were extracted from the arCOG database [arCOG14<sup>1</sup> (arCOG15272, arCOG00329, arCOG00328, arCOG04926, arCOG15270)]. Additional thaumarchaeotal *polB*

sequences were identified using “*Ca. N. islandicus*” as a query in a blastp search against the nr protein database. These additional thaumarchaeotal sequences, the *polB* sequence from “*Ca. N. islandicus*” and arCOG database sequences were de-replicated using usearch (Edgar, 2010) with -sortbylength and -cluster\_smallmem (-id 0.99 -query\_cov 0.9), aligned using default settings in mafft (Katoh and Standley, 2013) and a phylogenetic tree was calculated using FastTree (Price et al., 2010).

Nitrilase superfamily amino acid sequences were obtained from Pace and Brenner (2001). Alignment and phylogenetic reconstruction was carried out with Bali-Phy (Suchard and Redelings, 2006; randomize alignment, iterations = 11000, burnin = 6000). Posterior tree pools from 10 independent runs were combined to generate a 50% PP consensus tree and to assess bipartition support.

A dataset for assessing the phylogenetic relationship of the alpha subunit of 2-oxoacid:ferredoxin oxidoreductases (OFORs) was based on Gibson et al. (2016) and supplemented with additional indolepyruvate oxidoreductase (*ior*) sequences. Genomes available (as of October 30, 2017) from the NCBI genomes database were downloaded, genes were predicted using Prodigal V2.6.3 (Hyatt et al., 2010) and predicted genes were screened for *iorA* (TIGRFAM03336) using hmmsearch v3.1b2<sup>2</sup> with an *e*-value cutoff of 0.001. Genes meeting the search criteria were used as queries against the complete TIGRFAM database to ensure that the extracted *iorA* sequences matched the *iorA* model as the best-hit model with an *e*-value cutoff of 0.001. Reciprocal best-hit genes were required to align to the hmm over at least 500 contiguous bases. Amino acid sequences were then clustered into centroids using usearch v8.0.1517 (sortbylength and cluster\_smallmem -id 0.8 -query\_cov 0.9; Edgar, 2010). Centroids were aligned using mafft v7.245 (Katoh and Standley, 2013) and trees were constructed using FastTree 2.1.4 (Price et al., 2010). The initial phylogenetic placement of “*Ca. N. islandicus*” *iorA* in the resulting large tree (3,179 sequences) was used to choose a small set of bacterial *iorA* sequences to include in the final tree. The final dataset was aligned using mafft v7.245 (Katoh and Standley, 2013) and trees were constructed using FastTree 2.1.4 (Price et al., 2010).

## Electron Microscopy

For scanning electron microscopy, “*Ca. N. islandicus*” cells were harvested by centrifugation (4,500  $\times$  g, 15 min, 25°C) and fixed on poly-L-lysine coated slides with a filter-sterilized 2.5% glutaraldehyde fixation solution in phosphate buffered saline (PBS; 130 mM NaCl in 5% [v/v] phosphate buffer mixture [20–80 v/v] of 200 mM NaH<sub>2</sub>PO<sub>4</sub> and 200 mM Na<sub>2</sub>HPO<sub>4</sub>). Subsequently, fixed cells were washed three times for 10 min in PBS and post-fixed with a 1% OsO<sub>4</sub> solution in PBS for 40 min. The fixed cells were again washed three times in PBS, dehydrated in a 30–100% (v/v) ethanol series, washed in acetone, and critical point dried with a CPD 300 unit (Leica). Samples were mounted on stubs, sputter coated with gold using a sputter coater JFC-2300HR (JEOL), and images were obtained with a JSM-IT300 scanning electron microscope (JEOL).

<sup>1</sup> <ftp://ftp.ncbi.nih.gov/pub/wolf/COGs/arCOG/>

<sup>2</sup> <http://hmmer.org>


## RESULTS AND DISCUSSION

### Enrichment and Basic Physiology of “*Ca. N. islandicus*”

An ammonia-oxidizing enrichment culture was established from biofilm material sampled from a hot spring located in the geothermal valley Graendalur of South-Western Iceland. Temperature tests for optimal activity and growth were performed at different time points during the enrichment period and showed varying results, but below 50°C and above 75°C activity and growth was never observed. Only during the initial enrichment phase did ammonia oxidation occur at 75°C. At 65°C the highest ammonia oxidation rates and the shortest lag phases were usually measured (data not shown), however, in a single experiment the optimal temperature was 70°C (Supplementary Figure S1). Likely, these variations reflect varying abundance ratios of “*Ca. N. islandicus*” and accompanying bacteria over time as described in Lebedeva et al. (2008). A high enrichment level of a single AOA phylotype (see below) was achieved by applying the antibiotic spiramycin (15 mg l<sup>-1</sup>) followed by biomass transfers into fresh medium using serial dilutions. This enrichment culture showed near stoichiometric conversion of ammonium to nitrite when incubated at 65°C (Figure 1). This was accompanied by growth of the AOA with a specific growth rate of  $0.128 \pm 0.011 \text{ d}^{-1}$  (mean generation time of  $2.32 \pm 0.24 \text{ d}$ ), which is substantially slower than those reported for “*Ca. Nitrosocaldus yellowstonensis* HL72,” *Nitrososphaera viennensis* EN76, or *Nitrosopumilus maritimus* SCM1 (Könneke et al., 2005; de la Torre et al., 2008; Martens-Habben et al., 2009; Stieglmeier et al., 2014b; Table 1), but faster than a marine enrichment culture (Berg et al., 2015). At late exponential phase “*Ca. N. islandicus*” grown at 65°C reached a density of  $1.90 \times 10^4 \pm 2.84 \times 10^3$  cells per  $\mu\text{l}$ .

### Genome Reconstruction, Phylogeny, and Environmental Distribution

Metagenomic sequencing of the enrichment culture with Illumina and Nanopore demonstrated that the current culture contained an AOA as the only taxon encoding the repertoire genes required for ammonia oxidation. Hybrid assembly allowed reconstruction of the complete genome of this AOA as one circular contiguous sequence of 1.62 Mbps length (Table 1). The 16S rRNA gene and *amoA* gene of the newly enriched AOA are 96 and 85% identical, respectively, to the genes of “*Ca. Nitrosocaldus yellowstonensis*,” the only other cultured obligately thermophilic AOA. The average amino acid sequence identity (AAI) and the genomic average nucleotide identity (gANI) between the genome and the one of “*Ca. N. yellowstonensis*” are 65.4% (alignment fraction: 0.86) and 75.8% (alignment fraction: 0.59), respectively, which is above the proposed genus and below the proposed species boundary thresholds (Qin et al., 2014; Varghese et al., 2015). Consequently, the enriched obligately thermophilic AOA was assigned to the same genus and referred to as “*Ca. Nitrosocaldus islandicus*.” According to 16S rRNA gene- based phylogenies, “*Ca. N. islandicus*” is a member of the Nitrosocaldales clade, which seems to predominantly encompass AOA from thermal environments (Figure 2). An extended phylogenomic analysis using a concatenated alignment of 34 proteins (Supplementary Table S2) identified by CheckM (Parks et al., 2015) confirmed that “*Ca. N. islandicus*” represents a basal lineage within the known ammonia-oxidizing Thaumarchaeota (Figure 3). This result lends strong support to the earlier notion, which was based on single-gene 16S rRNA and *amoA* phylogenies (de la Torre et al., 2008), that the thermophilic Nitrosocaldales clade is an early diverging group of the ammonia-oxidizing Thaumarchaeota. It would also be compatible with the possibility that archaeal ammonia oxidation originated in thermal environments (de la Torre et al., 2008; Hatzenpichler



**FIGURE 1 |** Near-stoichiometric oxidation of 1.25 mM ammonium to nitrite by the “*Ca. N. islandicus*” enrichment culture at 65°C. Data points show means, error bars show 1 SD of  $n = 3$  biological replicates. Solid and dashed lines denote live and dead culture incubations, respectively. If not visible, error bars are smaller than symbols.

**TABLE 1** | Genome features and growth rates of “*Candidatus N. islandicus*” and of selected reference ammonia-oxidizing archaea (AOA).

Genome features	“ <i>Ca. N. islandicus</i> 3F”	<i>N. gargensis</i> Ga9-2	<i>N. viennensis</i> EN76	“ <i>Ca. N. exaquare</i> G61”	“ <i>Ca. N. devanattera</i> Nd1”	“ <i>Ca. N. uzonensis</i> N4”	<i>N. maritimus</i> SCM1
Genome size [Mb]	1.62	2.83	2.53	2.99	1.81	1.65	1.60
Number of scaffolds	1	1	1	1	1	1	1
Number of contigs	1	1	1	1	1	14	1
Average G+C content [%]	41.54	48.35	52.72	33.94	37.07	42.25	34.17
Protein coding density [%]	87.85	83.37	86.43	77.14	90.55	90.42	91.65
Number of genomic objects (CDS, fragment CDS, r/tRNA)	1851	4037	3266	3394	2145	2001	2012
Number of coding sequences (CDS)	1824	3999	3277	3358	2106	1960	1969
Motility/chemotaxis	+	+	+	–	+	+	–
Carbon fixation	3HP/4HB	3HP/4HB	3HP/4HB	3HP/4HB	3HP/4HB	3HP/4HB	3HP/4HB
Ammonium transporters	3	3	3	1	3	2	2
NirK	0	1	1	1	1	1	1
MCO1 + ZIP/MCO1 <sup>a</sup>	1/0	1/1	2/0	1/1	0/0	1/0	2/0
Urease and urea transport	+	+	+	+	–	–	–
Cyanate lyase	–	+	–	–	–	–	–
Nitrilase/Cyanide hydratase	1	0	0	0	0	1	1
Aromatic amino acid fermentation	+	–	–	–	–	–	–
Hydrogenase	3b	–	4a	–	4a	–	–
Coenzyme F420	+	+	+	+	+	+	+
Vitamin B12	+	+	+	+	+	+	+
Catalase	0	(1) <sup>b</sup>	0	1	0	0	0
Peroxidase	0	0	0	1	0	0	0
Superoxide dismutase	1	1	1	1	1	1	2
Chlorite dismutase-like enzyme <sup>c</sup>	1	1	1	1	1	1	1
DNA polymerases	B1, Y	B1, D, Y	B1, D, Y	B1, D, Y	B1, D, Y	B1, D, Y	B1, D, Y
Generation time [d]	2.32 ± 0.24 <sup>d</sup>	NA	1.25 ± 0.03	NA	NA	NA	0.88 – 1.08

<sup>a</sup>MCO1+ZIP/MCO1, multicopper oxidase 1 (as defined in Kerou et al., 2016) with adjacent zinc permease/multicopper oxidase 1 without an adjacent zinc permease;

<sup>b</sup>The gene is truncated; <sup>c</sup>chlorite dismutases are of interest in other nitrifiers (Maixner et al., 2008; Kostan et al., 2010), but it is not known what their function is in archaea;

<sup>d</sup>determined at 65°C; NA, not available.

et al., 2008; Groussin and Gouy, 2011; Brochier-Armanet et al., 2012).


Metagenomic sequencing revealed that in addition to “*Ca. N. islandicus*” the culture also contained two heterotrophic bacterial contaminants, which were identified as a *Thermus* sp. and a member of the Chloroflexi phylum (Figure 4). The enrichment level of “*Ca. N. islandicus*” was approximately 85% based on read counts from the Nanopore sequencing, whereas the *Thermus* sp. and Chloroflexi accounted for 12 and 3%, respectively. FISH-analysis of the enrichment culture confirmed the dominance of “*Ca. N. islandicus*” and showed that the AOA grew mainly in aggregates, whereas the bacterial cells grew either co-localized with the archaeal flocs or planktonically (Figure 5A). Electron microscopy demonstrated that the cells of “*Ca. N. islandicus*” are small (with a diameter of approximately 0.5–0.7 μm) and have an irregular coccoid shape (Figure 5B). Morphologically they resemble the cells of “*Ca. N. yellowstonensis*” (Qin et al., 2017b).

The environmental distribution of the two cultured Nitrosocaldales members and closely related AOA was assessed by screening all publicly available 16S rRNA gene amplicon datasets ( $n = 93,045$ ) for sequences highly similar (97%) to the 16S rRNA genes of “*Ca. N. islandicus*” and “*Ca. N. yellowstonensis*” using the pipeline described by Lagkouvardos

et al. (2016). This analysis revealed that these taxa are highly confined in their distribution and occur predominantly in terrestrial hot springs where they can reach high relative abundances between 11.4 and 86% (“*Ca. N. islandicus*” and “*Ca. N. yellowstonensis*,” respectively) of the total microbial community (Figure 6). Interestingly, “*Ca. N. yellowstonensis*”-related organisms seem to occur mainly in hot springs described as alkaline with a pH of around 8.5, but were also detected in a sample from a Tibetan wastewater treatment plant (Niu et al., 2017). The unexpected detection of members of the Nitrosocaldales in the latter sample was confirmed by 16S rRNA-based phylogenetic analyses (data not shown) and it would be interesting to know whether this wastewater treatment plant is in some way connected to water from a close-by hot spring.

## Genome Features


Addition of the complete genome of “*Ca. N. islandicus*” to the set of available thaumarchaeotal genome sequences ( $n = 30$ ) reduced the number of gene families identified as representing the “Thaumarchaeota-core” (Herbold et al., 2017) from 743 to 669 (reduction by 9.96%; Supplementary Table S3). In a few cases, genes with low sequence homology to apparently absent core gene families are actually present in the genome of “*Ca.*




**FIGURE 2 |** 16S rRNA gene-based (A) and *amoA* gene-based (B) maximum likelihood phylogenies of representative thaumarchaeotal sequences. For each sequence, the accession number and environmental source are indicated. Sequences from pure and enrichment cultures are depicted in bold, and “Ca. N. islandicus” is highlighted in red. The outgroup for the 16S rRNA tree were algarachaeal sequences; the *amoA* phylogeny was calculated unrooted, but artificially rooted to the Nitrosocaldus afterwards. Circles at nodes denote support (filled) or no support (open) from Neighbour Joining (NJ, top half) and Maximum Parsimony (MP, bottom half) trees. The scale bars in (A,B) indicate 9 and 6% sequence divergence, respectively.







**FIGURE 4 |** Sequence composition-independent binning of the metagenome scaffolds from two ammonia-oxidizing enrichment cultures. Circles represent scaffolds, scaled by the square root of their length. Clusters of similarly colored circles represent potential genome bins. The x-axis shows binning of the scaffolds from an early enrichment culture, which still included other genera as well (not shown). The y-axis shows binning of the scaffolds from the latest enrichment culture containing only “*Ca. N. islandicus*” and the two remaining accompanying organisms. Genome bins for the *Thermus* (34% complete) and the *Chloroflexi* (56% complete) organism were obtained. The genome bin of the *Chloroflexi* organism contains genes that cluster within a clade of *Nitrobacter/Nitrolancea* nitrite oxidoreductase (*nrxAB*) genes (data not shown). Some nitrate production by the batch enrichment culture was observed after multiple re-feedings with  $\text{NH}_4^+$  (>20 mM). As NOB in the phylum *Chloroflexi* are known (Sorokin et al., 2012), it is tempting to speculate that this *Chloroflexi* may be a thermophilic nitrite oxidizer.




**FIGURE 5 | (A)** FISH analysis of the enrichment culture illustrating the growth in microcolonies and the high relative abundance of “*Ca. N. islandicus*.” “*Ca. N. islandicus*” cells appear in green (stained by probe Thaum726 targeting most Thaumarchaeota) and the bacterial contaminants in blue (labeled by probe EUB338). **(B)** Scanning electron micrograph of spherically shaped “*Ca. Nitrosocaldus islandicus*” cells. The cells have a diameter of 0.5–0.7  $\mu\text{m}$ . “*Ca. N. islandicus*” cells were distinguishable from the rod-shaped bacterial contaminants by their smaller size and unique, ‘dented’ spherical shape.

In this context it is interesting to note that the hydroxylamine dehydrogenase of AOB, of which the functional homolog in archaea has not been identified yet, has recently been reported

to produce NO instead of nitrite (Caranto and Lancaster, 2017). Alternatively, NO could be provided by accompanying organisms such as the *Thermus* and *Chloroflexi*-like bacteria that remain in the enrichment. Indeed, the genome bins obtained for these organisms both encode a *nirK* gene. The *Thermus* sp. genome bin further contains a *norBC* and *narGH* genes, in line with described denitrification capabilities for the genus *Thermus* (Alvarez et al., 2014). A dependence of *Nitrosocaldus* on NO production by other microorganisms could explain why no pure culture from this lineage has been obtained yet.

“*Ca. N. islandicus*” possesses genes coding for urease that are present in some but not all AOA (Walker et al., 2010; Spang et al., 2012; Kerou et al., 2016; Lehtovirta-Morley et al., 2016; Sauder et al., 2017) (Figure 7 and Table 1), but lacks a cyanase that is used by *Nitrososphaera gargensis* for cyanate-based growth (Palatinszky et al., 2015). Additionally, the genome encodes an enzyme that either belongs to a novel class of the nitrilase superfamily or to the cyanide hydratase family (Figure 7 and Supplementary Figure S4). Nitrilases catalyze the direct cleavage of a nitrile to the corresponding acid while forming ammonia (Pace and Brenner, 2001) and cyanide hydratases convert HCN to formamide. Both substrates are relatively thermostable (Isidorov et al., 1992; Miyakawa et al., 2002). Nitriles occur as intermediates of microbial metabolism (Kobayashi et al., 1993) and nitrile hydratases have previously been isolated from several thermophiles (Cramp et al., 1997; Almatawah et al., 1999; Kabaivanova et al., 2008). Furthermore, both compounds are intermediates of the proposed abiotic synthesis of organics at hydrothermal sites (Miller and Urey, 1959; Schulte and Shock, 1995) and could thus be available in the hot spring habitat of “*Ca. N. islandicus*.” Similar genes have been found in the genomes of several other AOA from the *Nitrosopumilus* and *Nitrosotenuis* genera (Walker et al., 2010; Mosier et al., 2012; Lebedeva et al., 2013; Park et al., 2014; Bayer et al., 2016) (Table 1) and it will be interesting to find out for which metabolism they may be used in AOA.

Intriguingly, “*Ca. N. islandicus*” might be able to ferment amino acids under anaerobic conditions as it contains the entire pathway used by some hyperthermophilic archaea for ATP generation from aromatic amino acids (Mai and Adams, 1994; Adams et al., 2001; Ozawa et al., 2012) (Figure 7). In this pathway arylpyruvates are formed from aromatic amino acids by the activity of amino acid aminotransferases using 2-oxoglutarate as the amine group acceptor. The glutamate produced by this transamination can be recycled back to 2-oxoglutarate via glutamate dehydrogenase (*gdhA*) with the concomitant reduction of  $\text{NADP}^+$ . With *ilvE* and *aspC* genes present, “*Ca. N. islandicus*” encodes at least two enzymes for which an aminotransferase activity specific for tyrosine, phenylalanine and aspartate has been demonstrated (Gelfand and Steinberg, 1977). Subsequently, these 2-ketoacids could be oxidatively decarboxylated and converted to aryl-CoAs by the oxygen-sensitive enzyme indolepyruvate oxidoreductase (Ozawa et al., 2012) encoded by *iorAB* using oxidized ferredoxin as electron acceptor. *IorAB* is absent from all other genome-sequenced AOA and does also not occur in the ancestral thaumarchaeote “FN1” (Lin et al., 2015) that lacks AMO. The *ior* genes present in “*Ca. N. islandicus*” have




the highest similarity to and cluster together with *iorAB*-genes found in *Kyrpidia tusciae* and *Dadabacteria* (Supplementary Figure S5). Finally, transformation of aryl-CoAs to aryl acids catalyzed by ADP-dependent acetyl-CoA/acyl-CoA synthetase (Glasemacher et al., 1997) leads to ATP formation via substrate-level phosphorylation (Figure 7). “*Ca. N. islandicus*” encodes four acetyl-CoA/acyl-CoA synthetases, two of which are most similar to non-syntentous homologs of acetyl-CoA/acyl-CoA synthetases found in other AOA. However, the third gene is absent in all other AOA to date and its most similar homologs are encoded by species of the peptidolytic thermophilic archaea *Thermoproteus* and *Sulfolobus* and the fourth is most similar to an acetyl-/acyl-CoA synthetase found in members of the thermophilic Bathyarchaea and Hadesarchaea.

The fermentation of aromatic amino acids also requires regeneration of oxidized ferredoxin (reduced by *IorAB*) and NADP<sup>+</sup> (reduced by glutamate dehydrogenase). However, no canonical ferredoxin:NADP<sup>+</sup> oxidoreductase, or other enzymes (Buckel and Thauer, 2013) described to regenerate oxidized

ferredoxin, are encoded in the genome of “*Ca. N. islandicus*.” It seems unlikely that the amount of ferredoxin oxidized by an encoded ferredoxin-dependent assimilatory sulfite/nitrite reductase (Figure 7) would be sufficient to compensate for all ferredoxin reduced in the dissimilatory fermentation pathway. However, “*Ca. N. islandicus*” can also oxidize reduced ferredoxin with a 2-oxoglutarate-ferredoxin oxidoreductase (Supplementary Figure S5). NAD(P)H can be re-oxidized by a cytosolic, bidirectional, NAD(P)-coupled type 3b [NiFe] –hydrogenase that is encoded by “*Ca. N. islandicus*” in contrast to all other genomically characterized AOA (Figure 7 and Table 1). NAD(P)H oxidation by this hydrogenase could lead to hydrogen generation, or the enzyme could act as a sulfhydrogenase that reduces zero valent sulfur compounds (produced by other organisms or present in the environment) to hydrogen sulfide (Ma et al., 1993; Adams et al., 2001). The hydrogenase genes are clustered at a single locus and code for the four subunits of the holoenzyme and accessory proteins (Supplementary Figure S6). This hydrogenase might also allow “*Ca. N. islandicus*” to use hydrogen as energy source providing reduced NAD(P)H under oxic conditions as this type of hydrogenase has been shown to tolerate exposure to oxygen (Bryant and Adams, 1989; Berney et al., 2014; Kwan et al., 2015).

Surprisingly, the genome of “*Ca. N. islandicus*” lacks genes for both subunits of the DNA polymerase D (PolD), which is present in all other AOA and most archaeal lineages (including thermophiles) with the exception of the Crenarchaea (Cann et al., 1998; Makarova et al., 2014; Saw et al., 2015) (Table 1). It is assumed that either PolD alone or together with DNA polymerases of the B family (PolB) is required for DNA synthesis and elongation in these archaea (Cubonová et al., 2013; Ishino and Ishino, 2013; Makarova et al., 2014). The “*Ca. N. islandicus*” genome encodes only one B-type DNA polymerase (PolB1, Supplementary Figure S7) and one DNA polymerase of the Y family (PolY), generally considered to be involved in the rescue of stalled replication forks and enhancement of cell survival upon DNA damage (Friedberg et al., 2002). Recently, it has been demonstrated for the PolD-lacking crenarchaeon *Sulfolobus acidocaldarius* that both its PolB1 and PolY have polymerase activities *in vitro* (Peng et al., 2016). However, “*Ca. N. islandicus*” (like other AOA) does not encode the PolB1-binding proteins PBP1 and PBP2, which are required to form a multisubunit DNA polymerase holoenzyme together with PolB in the crenarchaeon *S. solfataricus* P2 (Yan et al., 2017). We hypothesize that “*Ca. N. islandicus*” may utilize one or both of the present polymerases for DNA replication, possibly in combination with its heterodimer PriSL, which has been demonstrated to function as a primase, a terminal transferase and a polymerase capable of polymerizing RNA or DNA chains of up to 7,000 nucleotides (Lao-Sirieix and Bell, 2004).

It is also interesting to note that the obligate thermophile “*Ca. N. islandicus*” like all genome-sequenced Thaumarchaeota (Spang et al., 2017) does not encode a reverse gyrase, which is widespread in hyperthermophilic microbes including other archaea of the TACK superphylum (Makarova et al., 2007; Heine and Chandra, 2009; López-García et al., 2015), but is not essential for growth under these conditions (Atomi et al., 2004).



## CONCLUSION

We have obtained a highly enriched (~85%) culture of an obligately thermophilic AOA from a hot spring in Iceland. Despite the impressive diversity of AOA in high temperature environments as revealed by molecular tools (Zhang et al.,

2008; Wang et al., 2009; Zhao et al., 2011; Nishizawa et al., 2013; Li et al., 2015; Chen et al., 2016), cultivation of only a single obligately thermophilic AOA species – “*Ca. Nitrosocaldus yellowstonensis*” – was reported before (de la Torre et al., 2008). The newly enriched AOA represents a new species of the genus *Nitrosocaldus* and was named “*Ca. N. islandicus*.” Comparative



analysis of its closed genome revealed several surprising features like the absence of DNA polymerase D and the lack of canonical NO-generating enzymes<sup>§</sup>, although physiological experiments with a NO-scavenger demonstrated NO-dependent ammonia-oxidation, as described for other AOA (Shen et al., 2013; Jung et al., 2014; Martens-Habben et al., 2015; Sauder et al., 2016). Furthermore, “*Ca. N. islandicus*” encodes the enzymatic repertoire for fermentation of aromatic amino acids that is, so far, unique among sequenced AOA. A pure culture of “*Ca. N. islandicus*” will be required to physiologically verify this genome-based hypothesis. Peptide or aromatic amino acid fermentation would enable an anaerobic lifestyle of “*Ca. N. islandicus*” and, if more widespread among Thaumarchaeota not yet characterized (including mesophiles), might help explain their sometimes surprisingly high abundance in anaerobic ecosystems (Molina et al., 2010; Bouskill et al., 2012; Buckles et al., 2013; Beam et al., 2014; Lin et al., 2015).

Based on the data presented here, we propose the following provisional taxonomic assignment for the thaumarchaeon in our enrichment culture.

Nitrosocaldales order

Nitrosocaldaceae fam.

‘*Candidatus Nitrosocaldus islandicus*’ sp. nov.

## Etymology

Nitrosus (Latin masculine adjective): nitrous; caldus (Latin masculine adjective): hot; islandicus (Latin masculine genitive name): from Iceland. The name alludes to the physiology of the organism (ammonia oxidizer, thermophilic) and the habitat from which it was recovered.

## Locality

The biofilm of a terrestrial hot spring in Graendalur geothermal valley, Iceland (64° 1’7’’ N, 21° 11’20’’ W).

## Diagnosis

An obligately thermophilic, aerobic chemolithoautotrophic ammonia oxidizer from the phylum Thaumarchaeota growing as small irregular shaped cocci. The values of AAI and gANI between this species and its closest cultured relative, “*Ca. N. yellowstonensis*”, are 65.4 and 75.8%, respectively.

## NOTE ADDED IN PROOF

After submission of this manuscript a second report on the enrichment and genome analysis of a thermophilic

<sup>§</sup> See section “Note Added in Proof”.

## REFERENCES

- Abell, G. C. J., Revell, A. T., Smith, C., Bissett, A. P., Volkman, J. K., and Robert, S. S. (2010). Archaeal ammonia oxidizers and nirS-type denitrifiers dominate sediment nitrifying and denitrifying populations in a subtropical macrotidal estuary. *ISME J.* 4, 286–300. doi: 10.1038/ismej.2009.105
- Adair, K. L., and Schwartz, E. (2008). Evidence that ammonia-oxidizing archaea are more abundant than ammonia-oxidizing bacteria in semiarid soils of

thaumarchaeote was submitted and accepted for publication in *Frontiers in Microbiology* (doi: 10.3389/fmicb.2018.00028). The genomes of both organisms share many of the above discussed features such as the absence of *nirK* and genes encoding polymerase D, and thereby lend support to the possibility that these are common genomic features of the Nitrosocaldales.

## AUTHOR CONTRIBUTIONS

AD, JV, and CS cultivated and enriched the culture. AD, CS, and PP performed growth and activity experiments. AD performed FISH and SEM analysis. CH, AD, PP, MA, and RK performed bioinformatic analysis. JdlT kindly provided access to the “*Ca. N. yellowstonensis*” genome. AD, JV, CS, PP, MW, and HD manually curated the annotation of the genome and interpreted the genome data. AD and MW wrote the manuscript with help from all co-authors.

## FUNDING

JV, CS, CH, and MW were supported by the ERC Advanced Grant NITRICARE (294343) to MW. AD and HD were supported by the Austrian Science Fund (FWF) grants P25231-B21 and T938. MA was supported by a research grant (15510) from VILLUM FONDEN.

## ACKNOWLEDGMENTS

We thank Prof. Bjarni Sigurdson and the Agricultural University of Iceland for granting us permission to access and sample in Graendalur Valley, Iceland. Daniela Gruber from the Core Facility of Cell Imaging and Ultrastructure Research at the University of Vienna and Stefano Romano are gratefully acknowledged for help with sample preparation for electron microscopy. We are grateful to Anja Spang for fruitful discussions.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2018.00193/full#supplementary-material>

northern arizona, USA. *Microb. Ecol.* 56, 420–426. doi: 10.1007/s00248-007-9360-9

- Adams, M. W. W., Holden, J. F., Menon, A. L., Schut, G. J., Grunden, A. M., Hou, C., et al. (2001). Key role for sulfur in peptide metabolism and in regulation of three hydrogenases in the hyperthermophilic archaeon *Pyrococcus furiosus*. *J. Bacteriol.* 183, 716–724. doi: 10.1128/JB.183.2.716-724.2001
- Almatawah, Q. A., Cramp, R., and Cowan, D. A. (1999). Characterization of an inducible nitrilase from a thermophilic bacillus. *Extremophiles* 3, 283–291. doi: 10.1007/s007920050129

- Alvarez, L., Bricio, C., Blesa, A., Hidalgo, A., and Berenguer, J. (2014). Transferable denitrification capability of *Thermus thermophilus*. *Appl. Environ. Microbiol.* 80, 19–28. doi: 10.1128/AEM.02594-13
- Amann, R. I., Binder, B. J., Olson, R. J., Chisholm, S. W., Devereux, R., and Stahl, D. A. (1990). Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations. *Appl. Environ. Microbiol.* 56, 1919–1925.
- Angel, R., and Conrad, R. (2013). Elucidating the microbial resuscitation cascade in biological soil crusts following a simulated rain event. *Environ. Microbiol.* 15, 2799–2815. doi: 10.1111/1462-2920.12140
- Atomi, H., Matsumi, R., and Imanaka, T. (2004). Reverse gyrase is not a prerequisite for hyperthermophilic life. *J. Bacteriol.* 186, 4829–4833. doi: 10.1128/JB.186.14.4829-4833.2004
- Bartossek, R., Nicol, G. W., Lanzen, A., Klenk, H. P., and Schleper, C. (2010). Homologues of nitrite reductases in ammonia-oxidizing archaea: diversity and genomic context. *Environ. Microbiol.* 12, 1075–1088. doi: 10.1111/j.1462-2920.2010.02153.x
- Bayer, B., Vojvoda, J., Offre, P., Alves, R. J. E., Elisabeth, N. H., Garcia, J. A., et al. (2016). Physiological and genomic characterization of two novel marine thaumarchaeal strains indicates niche differentiation. *ISME J.* 10, 1051–1063. doi: 10.1038/ismej.2015.200
- Beam, J. P. (2015). *Geobiological Interactions of Archaeal Populations in Acidic and Alkaline Geothermal Springs of Yellowstone National Park*. Bozeman, MT: Montana State University.
- Beam, J. P., Jay, Z. J., Kozubal, M. A., and Inskeep, W. P. (2014). Niche specialization of novel *Thaumarchaeota* to oxic and hypoxic acidic geothermal springs of Yellowstone National Park. *ISME J.* 8, 938–951. doi: 10.1038/ismej.2013.193
- Berg, C., Listmann, L., Vandieken, V., Vogts, A., and Jürgens K. (2015). Chemoautotrophic growth of ammonia-oxidizing *Thaumarchaeota* enriched from a pelagic redox gradient in the Baltic Sea. *Front. Microbiol.* 5:786. doi: 10.3389/fmicb.2014.00786
- Berney, M., Greening, C., Conrad, R., Jacobs, W. R., and Cook, G. M. (2014). An obligately aerobic soil bacterium activates fermentative hydrogen production to survive reductive stress during hypoxia. *Proc. Natl. Acad. Sci. U.S.A.* 111, 11479–11484. doi: 10.1073/pnas.1407034111
- Bouskill, N. J., Eveillard, D., Chien, D., Jayakumar, A., and Ward, B. B. (2012). Environmental factors determining ammonia-oxidizing organism distribution and diversity in marine environments. *Environ. Microbiol.* 14, 714–729. doi: 10.1111/j.1462-2920.2011.02623.x
- Brochier-Armanet, C., Boussau, B., Gribaldo, S., and Forterre, P. (2008). Mesophilic *Crenarchaeota*: proposal for a third archaeal phylum, the *Thaumarchaeota*. *Nat. Rev. Microbiol.* 6, 245–252. doi: 10.1038/nrmicro1852
- Brochier-Armanet, C., Gribaldo, S., and Forterre, P. (2012). Spotlight on the *Thaumarchaeota*. *ISME J.* 6, 227–230. doi: 10.1038/ismej.2011.145
- Bryant, F. O., and Adams, M. W. (1989). Characterization of hydrogenase from the hyperthermophilic archaeobacterium *Pyrococcus furiosus*. *J. Biol. Chem.* 264, 5070–5079.
- Buckel, W., and Thauer, R. K. (2013). Energy conservation via electron bifurcating ferredoxin reduction and proton/Na<sup>+</sup> translocating ferredoxin oxidation. *Biochim Biophys Acta* 1827, 94–113. doi: 10.1016/j.bbabi.2012.07.002
- Buckles, L. K., Villanueva, L., Weijers, J. W. H., Verschuren, D., and Damsté, J. S. S. (2013). Linking isoprenoidal GDGT membrane lipid distributions with gene abundances of ammonia-oxidizing *Thaumarchaeota* and uncultured crenarchaeotal groups in the water column of a tropical lake (Lake Challa, East Africa). *Environ. Microbiol.* 15, 2445–2462. doi: 10.1111/1462-2920.12118
- Cann, I. K. O., Komori, K., Toh, H., Kanai, S., and Ishino, Y. (1998). A heterodimeric DNA polymerase: evidence that members of *Euryarchaeota* possess a distinct DNA polymerase. *Proc. Natl. Acad. Sci. U.S.A.* 95, 14250–14255. doi: 10.1073/pnas.95.24.14250
- Caranto, J. D., and Lancaster, K. M. (2017). Nitric oxide is an obligate bacterial nitrification intermediate produced by hydroxylamine oxidoreductase. *Proc. Natl. Acad. Sci. U.S.A.* 114, 8217–8222. doi: 10.1073/pnas.1704504114
- Chen, S., Peng, X., Xu, H., and Ta, K. (2016). Nitrification of archaeal ammonia oxidizers in a high-temperature hot spring. *Biogeosciences* 13, 2051–2060. doi: 10.5194/bg-13-2051-2016
- Cramp, R., Gilmour, M., and Cowan, D. A. (1997). Novel thermophilic bacteria producing nitrile-degrading enzymes. *Microbiology* 143, 2313–2320. doi: 10.1099/00221287-143-7-2313
- Cubonová, L., Richardson, T., Burkhart, B. W., Kelman, Z., Connolly, B. A., Reeve, J. N., et al. (2013). Archaeal DNA polymerase D but not DNA polymerase B is required for genome replication in *Thermococcus kodakarensis*. *J. Bacteriol.* 195, 2322–2328. doi: 10.1128/JB.02037-12
- Daebeler, A., Abell, G. C. J., Bodelier, P. L. E., Bodrossy, L., Frampton, D. M. F., Hefting, M. M., et al. (2012). Archaeal dominated ammonia-oxidizing communities in Icelandic grassland soils are moderately affected by long-term N fertilization and geothermal heating. *Front. Microbiol.* 3:352. doi: 10.3389/fmicb.2012.00352
- Daims, H., Brühl, A., Amann, R., Schleifer, K.-H., and Wagner, M. (1999). The domain-specific probe EUB338 is insufficient for the detection of all bacteria: Development and evaluation of a more comprehensive probe Set. *Syst. Appl. Microbiol.* 22, 434–444. doi: 10.1016/S0723-2020(99)80053-8
- Daims, H., Stoecker, K., and Wagner, M. (2005). “Fluorescence in situ hybridisation for the detection of prokaryotes,” in *Molecular Microbial Ecology*, eds A. M. Osborn, and C. J. Smith (Abington: Taylor & Francis), 213–239.
- de la Torre, J. R., de la Christopher, B. W., Anitra, E. I., Martin, K., and David, A. S. (2008). Cultivation of a thermophilic ammonia oxidizing archaeon synthesizing crenarchaeol. *Environ. Microbiol.* 10, 810–818. doi: 10.1111/j.1462-2920.2007.01506.x
- Dodsworth, J. A., Hungate, B. A., and Hedlund, B. P. (2011). Ammonia oxidation, denitrification and dissimilatory nitrate reduction to ammonium in two US Great Basin hot springs with abundant ammonia-oxidizing archaea. *Environ. Microbiol.* 13, 2371–2386. doi: 10.1111/j.1462-2920.2011.02508.x
- Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26, 2460–2461. doi: 10.1093/bioinformatics/btq461
- Erguder, T. H., Nico, B., Lieven, W., Massimo, M., and Willy, V. (2009). Environmental factors shaping the ecological niches of ammonia-oxidizing archaea. *FEMS Microbiol. Rev.* 33, 855–869. doi: 10.1111/j.1574-6976.2009.00179.x
- Francis, C. A., Beman, J. M., and Kuypers, M. M. M. (2007). New processes and players in the nitrogen cycle: The microbial ecology of anaerobic and archaeal ammonia oxidation. *ISME J.* 1, 19–27. doi: 10.1038/ismej.2007.8
- Francis, C. A., Roberts, K. J., Beman, J. M., Santoro, A. E., and Oakley, B. B. (2005). Ubiquity and diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean. *Proc. Natl. Acad. Sci. U.S.A.* 102, 14683–14688. doi: 10.1073/pnas.0506625102
- Friedberg, E. C., Wagner, R., and Radman, M. (2002). Specialized DNA polymerases, cellular survival, and the genesis of mutations. *Science* 296, 1627–1630. doi: 10.1126/science.1070236
- Gelfand, D. H., and Steinberg, R. A. (1977). *Escherichia coli* mutants deficient in the aspartate and aromatic amino acid aminotransferases. *J. Bacteriol.* 130, 429–440.
- Gibson, M., Chen, P.-T., and Drennan, C. L. (2016). A structural phylogeny for understanding 2-oxoacid oxidoreductase function. *Curr. Opin. Struct. Biol.* 41, 54–61. doi: 10.1016/j.sbi.2016.05.011
- Glasmacher, J., Bock, A.-K., Schmid, R., and Schönheit, P. (1997). Purification and properties of acetyl-CoA synthetase (ADP-forming), an archaeal enzyme of acetate formation and ATP synthesis, from the hyperthermophile *Pyrococcus furiosus*. *Eur. J. Biochem.* 244, 561–567. doi: 10.1111/j.1432-1033.1997.00561.x
- Groussin, M., and Gouy, M. (2011). Adaptation to environmental temperature is a major determinant of molecular evolutionary rates in Archaea. *Mol. Biol. Evol.* 28, 2661–2674. doi: 10.1093/molbev/msr098
- Guy, L., and Ettema, T. J. G. (2011). The archaeal “TACK” superphylum and the origin of eukaryotes. *Trends Microbiol.* 19, 580–587. doi: 10.1016/j.tim.2011.09.002
- Hallam, S. J., Konstantinidis, K. T., Putnam, N., Schleper, C., Watanabe, Y., Sugahara, J., et al. (2006). Genomic analysis of the uncultivated marine crenarchaeote *Cenarchaeum symbiosum*. *Proc. Natl. Acad. Sci. U.S.A.* 103, 18296–18301. doi: 10.1073/pnas.0608549103
- Hatzenpichler, R., Lebedeva, E. V., Spieck, E., Stoecker, K., Richter, A., Daims, H., et al. (2008). A moderately thermophilic ammonia-oxidizing crenarchaeote from a hot spring. *Proc. Natl. Acad. Sci. U.S.A.* 105, 2134–2139. doi: 10.1073/pnas.0708857105

- Heine, M., and Chandra, S. B. C. (2009). The linkage between reverse gyrase and hyperthermophiles: a review of their invariable association. *J. Microbiol.* 47, 229–234. doi: 10.1007/s12275-009-0019-8
- Herbold, C. W., Lehtovirta-Morley, L. E., Jung, M.-Y., Jehmlich, N., Hausmann, B., Han, P., et al. (2017). Ammonia-oxidizing archaea living at low pH: Insights from comparative genomics. *Environ. Microbiol.* 19, 4939–4952. doi: 10.1111/1462-2920.13971
- Hyatt, D., Chen, G.-L., Locascio, P. F., Land, M. L., Larimer, F. W., and Hauser, L. J. (2010). Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. doi: 10.1186/1471-2105-11-119
- Ishino, Y., and Ishino, S. (2013). “DNA Replication in Archaea, the Third Domain of Life,” in *The Mechanisms of DNA Replication*, ed. S. David (Rijeka: InTech).
- Isidorov, V., Zenkevich, I., and Karpov, G. (1992). Volatile organic compounds in steam-gas outflows of several volcanoes and hydrothermal systems in Kamchatka. *J. Volc. Seis.* 13, 287–293.
- Jiang, H. C., Huang, Q. Y., Dong, H. L., Wang, P., Wang, F. P., Li, W. J., et al. (2010). RNA-based investigation of ammonia-oxidizing archaea in hot springs of Yunnan Province, China. *Appl. Environ. Microbiol.* 76, 4538–4541. doi: 10.1128/AEM.00143-10
- Jung, M.-Y., Park, S.-J., Kim, S.-J., Kim, J.-G., Damsté, J. S. S., Jeon, C. O., et al. (2014). A Mesophilic, autotrophic, ammonia-oxidizing archaeon of thaumarchaeal Group I.1a cultivated from a deep oligotrophic soil horizon. *Appl. Environ. Microbiol.* 80, 3645–3655. doi: 10.1128/AEM.03730-13
- Kabaivanova, L., Dimitrov, P., Boyadzhieva, I., Engibarov, S., Dobрева, E., and Emanuilova E. (2008). Nitrile degradation by free and immobilized cells of the thermophile *Bacillus* sp. UG-5B, isolated from polluted industrial waters. *World J. Microbiol. Biotechnol.* 24, 2383–2388. doi: 10.1007/s11274-008-9757-8
- Käll, L., Krogh, A., and Sonnhammer, E. L. L. (2007). Advantages of combined transmembrane topology and signal peptide prediction—the Phobius web server. *Nucleic Acids Res.* 35, W429–W432. doi: 10.1093/nar/gkm256
- Kandeler, E., and Gerber, H. (1988). Short-term assay of soil urease activity using colorimetric determination of ammonium. *Biol. Fertil. Soils* 6, 68–72. doi: 10.1007/BF00257924
- Katoh, K., and Standley, D. M. (2013). MAFFT Multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30, 772–780. doi: 10.1093/molbev/mst010
- Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N., and Sternberg, M. J. E. (2015). The Phyre2 web portal for protein modeling, prediction and analysis. *Nat. Protoc.* 10, 845–858. doi: 10.1038/nprot.2015.053
- Kerou, M., Offire, P., Valledor, L., Abby, S. S., Melcher, M., Nagler, M., et al. (2016). Proteomics and comparative genomics of *Nitrososphaera viennensis* reveal the core genome and adaptations of archaeal ammonia oxidizers. *Proc. Natl. Acad. Sci. U.S.A.* 113, E7937–E7946.
- Kobayashi, M., Izui, H., Nagasawa, T., and Yamada, H. (1993). Nitrilase in biosynthesis of the plant hormone indole-3-acetic acid from indole-3-acetonitrile: cloning of the *Alcaligenes* gene and site-directed mutagenesis of cysteine residues. *Proc. Natl. Acad. Sci. U.S.A.* 90, 247–251.
- Koch, H., Lückner, S., Albertsen, M., Kitzinger, K., Herbold, C., Spieck, E., et al. (2015). Expanded metabolic versatility of ubiquitous nitrite-oxidizing bacteria from the genus *Nitrospira*. *Proc. Natl. Acad. Sci. U.S.A.* 112, 11371–11376. doi: 10.1073/pnas.1506533112
- Könneke, M., Bernhard, A. E., de la Torre, J. R., Walker, C. B., Waterbury, J. B., and Stahl, D. A. (2005). Isolation of an autotrophic ammonia-oxidizing marine archaeon. *Nature* 437, 543–546. doi: 10.1038/nature03911
- Könneke, M., Schubert, D. M., Brown, P. C., Hügl, M., Standfest, S., Schwander, T., et al. (2014). Ammonia-oxidizing archaea use the most energy-efficient aerobic pathway for CO<sub>2</sub> fixation. *Proc. Natl. Acad. Sci. U.S.A.* 111, 8239–8244. doi: 10.1073/pnas.1402028111
- Konstantinidis, K. T., and Tiedje, J. M. (2005). Towards a genome-based taxonomy for prokaryotes. *J. Bacteriol.* 187, 6258–6264. doi: 10.1128/JB.187.18.6258-6264.2005
- Kostan, J., Sjöblom, B., Maixner, F., Mlynek, G., Furtmüller, P. G., Obinger, C., et al. (2010). Structural and functional characterisation of the chlorite dismutase from the nitrite-oxidizing bacterium “*Candidatus Nitrospira defluvi*”: identification of a catalytically important amino acid residue. *J. Struct. Biol.* 172, 331–342. doi: 10.1016/j.jsb.2010.06.014
- Kozłowski, J. A., Stieglmeier, M., Schleper, C., Klotz, M. G., and Stein, L. Y. (2016). Pathways and key intermediates required for obligate aerobic ammonia-dependent chemolithotrophy in bacteria and Thaumarchaeota. *ISME J.* 10, 1836–1845. doi: 10.1038/ismej.2016.2
- Kwan, P., McIntosh, C. L., Jennings, D. P., Hopkins, R. C., Chandrayan, S. K., Wu, C.-H., et al. (2015). The [NiFe]-hydrogenase of *Pyrococcus furiosus* exhibits a new type of oxygen tolerance. *J. Am. Chem. Soc.* 137, 13556–13565. doi: 10.1021/jacs.5b07680
- Lagkouvardos, I., Joseph, D., Kapfhammer, M., Girtli, S., Horn, M., Haller, D., et al. (2016). IMNGS: a comprehensive open resource of processed 16S rRNA microbial profiles for ecology and diversity studies. *Sci. Rep.* 6:33721. doi: 10.1021/srep33721
- Lao-Sirieix, S., and Bell, S. D. (2004). The heterodimeric primase of the hyperthermophilic archaeon *Sulfolobus solfataricus* possesses DNA and RNA primase, polymerase and 3'-terminal nucleotidyl transferase activities. *J. Mol. Biol.* 344, 1251–1263. doi: 10.1016/j.jmb.2004.10.018
- Lartillot, N., Lepage, T., and Blanquart, S. (2009). PhyloBayes 3: a Bayesian software package for phylogenetic reconstruction and molecular dating. *Bioinformatics* 25, 2286–2288. doi: 10.1093/bioinformatics/btp368
- Lebedeva, E. V., Alawi, M., Maixner, F., Jozsa, P.-G., Daims, H., and Spieck, E. (2008). Physiological and phylogenetic characterization of a novel lithoautotrophic nitrite-oxidizing bacterium, “*Candidatus Nitrospira bockiana*”. *Int. J. Syst. Evol. Microbiol.* 58, 242–250. doi: 10.1099/ijs.0.65379-0
- Lebedeva, E. V., Hatzepichler, R., Pelletier, E., Schuster, N., Hauzmayer, S., Bulaev, A., et al. (2013). Enrichment and genome sequence of the group I.1a ammonia-oxidizing archaeon “*Ca. Nitrosotenuis uzonensis*” representing a clade globally distributed in thermal habitats. *PLOS ONE* 8:e80835. doi: 10.1371/journal.pone.0080835
- Lehtovirta-Morley, L. E., Sayavedra-Soto, L. A., Gallois, N., Schouten, S., Stein, L. Y., Prosser, J. I., et al. (2016). Identifying potential mechanisms enabling acidophily in the ammonia-oxidizing archaeon “*Candidatus Nitrosotalea devanateri*”. *Appl. Environ. Microbiol.* 82, 2608–2619. doi: 10.1128/AEM.04031-15
- Leininger, S., Urich, T., Schloter, M., Schwark, L., Qi, J., Nicol, G. W., et al. (2006). Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature* 442, 806–809. doi: 10.1038/nature04983
- Li, H. (2016). Minimap and miniasm: fast mapping and de novo assembly for noisy long sequences. *Bioinformatics* 32, 2103–2110. doi: 10.1093/bioinformatics/btw152
- Li, H., Yang, Q., Li, J., Gao, H., Li, P., and Zhou, H. (2015). The impact of temperature on microbial diversity and AOA activity in the Tengchong geothermal field, China. *Sci. Rep.* 5:17056. doi: 10.1038/srep17056
- Lin, X., Handley, K. M., Gilbert, J. A., and Kostka, J. E. (2015). Metabolic potential of fatty acid oxidation and anaerobic respiration by abundant members of *Thaumarchaeota* and *Thermoplasmata* in deep anoxic peat. *ISME J.* 9, 2740–2744. doi: 10.1038/ismej.2015.77
- López-García, P., Zivanovic, Y., Deschamps, P., and Moreira, D. (2015). Bacterial gene import and mesophilic adaptation in archaea. *Nat. Rev. Microbiol.* 13, 447–456. doi: 10.1038/nrmicro3485
- Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadhukumar, et al. (2004). ARB: a software environment for sequence data. *Nucleic Acids Res.* 32, 1363–1371. doi: 10.1093/nar/gkh293
- Ma, K., Schicho, R. N., Kelly, R. M., and Adams, M. W. (1993). Hydrogenase of the hyperthermophile *Pyrococcus furiosus* is an elemental sulfur reductase or sulfhydrogenase: evidence for a sulfur-reducing hydrogenase ancestor. *Proc. Natl. Acad. Sci. U.S.A.* 90, 5341–5344. doi: 10.1073/pnas.90.11.5341
- Mai, X., and Adams, M. W. (1994). Indolepyruvate ferredoxin oxidoreductase from the hyperthermophilic archaeon *Pyrococcus furiosus*. A new enzyme involved in peptide fermentation. *J. Biol. Chem.* 269, 16726–16732.
- Maixner, F., Wagner, M., Lückner, S., Pelletier, E., Schmitz-Esser, S., Hae, K., et al. (2008). Environmental genomics reveals a functional chlorite dismutase in the nitrite-oxidizing bacterium “*Candidatus Nitrospira defluvi*”. *Environ. Microbiol.* 10, 3043–3056. doi: 10.1111/j.1462-2920.2008.01646.x
- Makarova, K. S., Krupovic, M., and Koonin, E. V. (2014). Evolution of replicative DNA polymerases in archaea and their contributions to the eukaryotic replication machinery. *Front. Microbiol.* 5:354. doi: 10.3389/fmicb.2014.00354



- Makarov, K. S., Sorokin, A. V., Novichkov, P. S., Wolf, Y. I., and Koonin, E. V. (2007). Clusters of orthologous genes for 41 archaeal genomes and implications for evolutionary genomics of archaea. *Biol. Direct* 2:33. doi: 10.1186/1745-6150-2-33
- Martens-Habbena, W., Berube, P. M., Urakawa, H., de la Torre, J. R., and Stahl, D. A. (2009). Ammonia oxidation kinetics determine niche separation of nitrifying archaea and bacteria. *Nature* 461, 976–979. doi: 10.1038/nature08465
- Martens-Habbena, W., Qin, W., Horak, R. E. A., Urakawa, H., Schauer, A. J., Moffett, J. W., et al. (2015). The production of nitric oxide by marine ammonia-oxidizing archaea and inhibition of archaeal ammonia oxidation by a nitric oxide scavenger. *Environ. Microbiol.* 17, 2261–2274. doi: 10.1111/1462-2920.12677
- Miller, S. L., and Urey, H. C. (1959). Organic compound synthesis on the primitive earth. *Science* 130, 245–251. doi: 10.1126/science.130.3370.245
- Mincer, T. J., Church, M. J., Taylor, L. T., Preston, C., Kar, D. M., and DeLong, E. F. (2007). Quantitative distribution of presumptive archaeal and bacterial nitrifiers in Monterey Bay and the North Pacific Subtropical Gyre. *Environ. Microbiol.* 9, 1162–1175. doi: 10.1111/j.1462-2920.2007.01239.x
- Miranda, K. M., Espey, M. G., and Wink, D. A. (2001). A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide* 5, 62–71. doi: 10.1006/niox.2000.0319
- Miyakawa, S., Cleaves, H. J., and Miller, S. L. (2002). The cold origin of life: Implications based on the hydrolytic stabilities of hydrogen, cyanide and formamide. *Orig. Life Evol. Biosph.* 32, 195–208. doi: 10.1023/A:1016514305984
- Molina, V., Belmar, L., and Ulloa, O. (2010). High diversity of ammonia-oxidizing archaea in permanent and seasonal oxygen-deficient waters of the eastern South Pacific. *Environ. Microbiol.* 12, 2450–2465. doi: 10.1111/j.1462-2920.2010.02218.x
- Mosier, A. C., Allen, E. E., Kim, M., Ferreira, S., and Francis, C. A. (2012). Genome Sequence of “*Candidatus Nitrosoarchaeum limnia*” BG20, a low-salinity ammonia-oxidizing archaeon from the San Francisco Bay estuary. *J. Bacteriol.* 194, 2119–2120. doi: 10.1128/JB.00007-12
- Musmann, M., Brito, I., Pitcher, A., Damste, J. S. S., Hatzenpichler, R., Richter, A., et al. (2011). Thaumarchaeotes abundant in refinery nitrifying sludges express *amoA* but are not obligate autotrophic ammonia oxidizers. *Proc. Natl. Acad. Sci. U.S.A.* 108, 16771–16776. doi: 10.1073/pnas.1106427108
- Nishizawa, M., Koba, K., Makabe, A., Yoshida, N., Kaneko, M., Hirao, S., et al. (2013). Nitrification-driven forms of nitrogen metabolism in microbial mat communities thriving along an ammonium-enriched subsurface geothermal stream. *Geochim. Cosmochim. Acta* 113, 152–173. doi: 10.1016/j.gca.2013.03.027
- Niu, L., Zhang, X., Li, Y., Wang, P., Zhang, W., Wang, C., et al. (2017). Elevational characteristics of the archaeal community in full-scale activated sludge wastewater treatment plants at a 3660-meter elevational scale. *Water Sci. Technol.* 76, 531–541. doi: 10.2166/wst.2017.215
- Otte, J., Mall, A., Schubert, D. M., Könneke, M., and Berg, I. A. (2015). Malonic semialdehyde reductase from the archaeon *Nitrosopumilus maritimus* is involved in the autotrophic 3-hydroxypropionate/4-hydroxybutyrate cycle. *Appl. Environ. Microbiol.* 81, 1700–1707. doi: 10.1128/AEM.03390-14
- Ozawa, Y., Siddiqui, M. A., Takahashi, Y., Urushiyama, A., Ohmori, D., Yamakura, F., et al. (2012). Indolepyruvate ferredoxin oxidoreductase: an oxygen-sensitive iron-sulfur enzyme from the hyperthermophilic archaeon *Thermococcus profundus*. *J. Biosci. Bioeng.* 114, 23–27. doi: 10.1016/j.jbiosc.2012.02.014
- Pace, H. C., and Brenner, C. (2001). The nitrilase superfamily: classification, structure and function. *Genome Biol.* 2, reviews0001.1–reviews0001.9. doi: 10.1186/gb-2001-2-1-reviews0001
- Palatinszky, M., Herbold, C., Jehmlich, N., Pogoda, M., Han, P., von Bergen, M., et al. (2015). Cyanate as an energy source for nitrifiers. *Nature* 524, 105–108. doi: 10.1038/nature14856
- Park, S.-J., Ghai, R., Martín-Cuadrado, A.-B., Rodríguez-Valera, F., Chung, W.-H., Kwon, K., et al. (2014). Genomes of two new ammonia-oxidizing archaea enriched from deep marine sediments. *PLOS ONE* 9:e96449. doi: 10.1371/journal.pone.0096449
- Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., and Tyson, G. W. (2015). CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* 25, 1043–1055. doi: 10.1101/gr.186072.114
- Peng, L., Xia, X., and Liu, X. (2016). Biochemical characterization of translesion synthesis by *Sulfolobus acidocaldarius* DNA polymerases. *Chem. Res. Chin. Univ.* 32, 226–233. doi: 10.1007/s40242-016-5337-x
- Pjevac, P., Schaubberger, C., Poghosyan, L., Herbold, C. W., van Kessel, M. A. H. J., Daebeler, A., et al. (2017). *AmoA*-targeted polymerase chain reaction primers for the specific detection and quantification of Comammox *Nitrospira* in the environment. *Front. Microbiol.* 8:1508. doi: 10.3389/fmicb.2017.01508
- Price, M. N., Dehal, P. S., and Arkin, A. P. (2010). FastTree 2 – Approximately maximum-likelihood trees for large alignments. *PLOS ONE* 5:e9490. doi: 10.1371/journal.pone.0009490
- Prosser, J. I., and Nicol, G. W. (2012). Archaeal and bacterial ammonia-oxidisers in soil: the quest for niche specialisation and differentiation. *Trends Microbiol.* 20, 523–531. doi: 10.1016/j.tim.2012.08.001
- Qin, Q.-L., Xie, B.-B., Zhang, X.-Y., Chen, X.-L., Zhou, B.-C., Zhou, J., et al. (2014). A proposed genus boundary for the prokaryotes based on genomic insights. *J. Bacteriol.* 196, 2210–2215. doi: 10.1128/JB.01688-14
- Qin, W., Heal, K. R., Ramdasi, R., Kobelt, J. N., Martens-Habbena, W., Bertagnolli, A. D., et al. (2017a). *Nitrosopumilus maritimus* gen. nov., sp. nov., *Nitrosopumilus cobalaminigenes* sp. nov., *Nitrosopumilus oxyclineae* sp. nov., and *Nitrosopumilus ureiphilus* sp. nov., four marine ammonia-oxidizing archaea of the phylum Thaumarchaeota. *Int. J. Syst. Evol. Microbiol.* doi: 10.1099/ijsem.0.002416 [Epub ahead of print].
- Qin, W., Jewell, T. N. M., Russell, V. V., Hedlund, B. P., de la Torre, J. R., and Stahl, D. A. (2017b). “*Candidatus Nitrosocaldus*,” in *Bergey’s Manual of Systematics of Archaea and Bacteria*, eds W. D. Grant, M. Kamekura, T. J. McGenity, and A. Ventosa (Hoboken, NJ: John Wiley & Sons).
- Reigstad, L. J., Richter, A., Daims, H., Urich, T., Schwark, L., and Schleper, C. (2008). Nitrification in terrestrial hot springs of Iceland and Kamchatka. *FEMS Microbiol. Ecol.* 64, 167–174. doi: 10.1111/j.1574-6941.2008.00466.x
- Sauder, L. A., Albertsen, M., Engel, K., Schwarz, J., Nielsen, P. H., Wagner, M., et al. (2017). Cultivation and characterization of *Candidatus Nitrosocosmicus exaquare*, an ammonia-oxidizing archaeon from a municipal wastewater treatment system. *ISME J.* 11, 1142–1157. doi: 10.1038/ismej.2016.192
- Sauder, L. A., Ross, A. A., and Neufeld, J. D. (2016). Nitric oxide scavengers differentially inhibit ammonia oxidation in ammonia-oxidizing archaea and bacteria. *FEMS Microbiol. Lett.* 363:fnw052. doi: 10.1093/femsle/fnw052
- Saw, J. H., Spang, A., Zaremba-Niedzwiedzka, K., Juzokaite, L., Dodsworth, J. A., Murugapiran, S. K., et al. (2015). Exploring microbial dark matter to resolve the deep archaeal ancestry of eukaryotes. *Philos. Trans. R. Soc. B* 370:20140328. doi: 10.1098/rstb.2014.0328
- Schleper, C., and Nicol, G. W. (2010). “Ammonia-oxidizing archaea - physiology, ecology, evolution,” in *Advances in Microbial Physiology*, Vol. 57, ed. R. K. Poole (London: Elsevier Ltd).
- Schulte, M., and Shock, E. (1995). Thermodynamics of strecker synthesis in hydrothermal systems. *Orig. Life Evol. Biosph.* 25, 161–173. doi: 10.1007/BF01581580
- Shen, T., Stieglmeier, M., Dai, J., Urich, T., and Schleper, C. (2013). Responses of the terrestrial ammonia oxidizing archaeon *Ca. Nitrososphaera viennensis* and the ammonia oxidizing bacterium *Nitrospira multififormis* to nitrification inhibitors. *FEMS Microbiol. Lett.* 344, 121–129. doi: 10.1111/1574-6968.12164
- Sorokin, D. Y., Lucker, S., Vejmolkova, D., Kostrikina, N. A., Kleerebezem, R., Rijpstra, W. I. C., et al. (2012). Nitrification expanded: discovery, physiology and genomics of a nitrite-oxidizing bacterium from the phylum Chloroflexi. *ISME J.* 6, 2245–2256. doi: 10.1038/ismej.2012.70
- Spang, A., Caceres, E. F., and Ettema, T. J. G. (2017). Genomic exploration of the diversity, ecology, and evolution of the archaeal domain of life. *Science* 357:eaa3883. doi: 10.1126/science.aaf3883
- Spang, A., Poehlein, A., Offre, P., Zumbragel, S., Haider, S., Rychlik, N., et al. (2012). The genome of the ammonia-oxidizing *Candidatus Nitrososphaera gargensis*: insights into metabolic versatility and environmental adaptations. *Environ. Microbiol.* 14, 3122–3145. doi: 10.1111/j.1462-2920.2012.02893.x
- Stahl, D. A., and Amann, R. (1991). “Development and application of nucleic acid probes in bacterial systematics,” in *Nucleic Acid Techniques in Bacterial Systematics*, eds E. Stackebrandt, and M. Goodfellow (Chichester: John Wiley & Sons Ltd.), 205–248.



- Stahl, D. A., and de la Torre, J. R. (2012). "Physiology and diversity of ammonia-oxidizing archaea," in *Annual Review of Microbiology*, Vol. 66 eds S. Gottesman, C. S. Harwood, and O. Schneewind (Palo Alto, CA: Annual Reviews, Inc.), 83–101.
- Stetter, K. O. (1998). "Hyperthermophiles: isolation, classification, and properties," in *Extremophiles: Microbial Life in Extreme Environments*, eds K. Horikoshi, and W. D. Grant (New York, NY: Wiley-Liss), 1–24.
- Stieglmeier, M., Alves, R. J. E., and Schleper, C. (2014a). "The phylum *Thaumarchaeota*," in *The Prokaryotes*, eds E. Rosenberg, E. F. DeLong, S. Lory, E. Stackebrandt, and F. Thompson (Berlin: Springer), 347–362.
- Stieglmeier, M., Klingl, A., Alves, R. J. E., Rittmann, S. K., Melcher, M., Leisch, N., et al. (2014b). *Nitrososphaera viennensis* sp. nov., an aerobic and mesophilic ammonia-oxidizing archaeon from soil and member of the archaeal phylum *Thaumarchaeota*. *Int. J. Syst. Evol. Microbiol.* 64, 2738–2752. doi: 10.1099/ijs.0.063172-0
- Suchard, M. A., and Redelings, B. D. (2006). BALI-Phy: simultaneous Bayesian inference of alignment and phylogeny. *Bioinformatics* 22, 2047–2048. doi: 10.1093/bioinformatics/btl175
- Tourna, M., Freitag, T. E., Nicol, G. W., and Prosser, J. I. (2008). Growth, activity and temperature responses of ammonia-oxidizing archaea and bacteria in soil microcosms. *Environ. Microbiol.* 10, 1357–1364. doi: 10.1111/j.1462-2920.2007.01563.x
- Vallenet, D., Belda, E., Calteau, A., Cruveiller, S., Engelen, S., Lajus, A., et al. (2013). MicroScope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucleic Acids Res.* 41, D636–D647. doi: 10.1093/nar/gks1194
- Varghese, N. J., Mukherjee, S., Ivanova, N., Konstantinidis, K. T., Mavrommatis, K., Kyrpides, N. C., et al. (2015). Microbial species delineation using whole genome sequences. *Nucleic Acids Res.* 43, 6761–6771. doi: 10.1093/nar/gkv657
- Vaser, R., Sović, I., Nagarajan, N., and Šikić, M. (2017). Fast and accurate de novo genome assembly from long uncorrected reads. *Genome Res.* 27, 737–746. doi: 10.1101/gr.214270.116
- Walker, C. B., Torre, J. R., de la Torre, M. G., Urakawa, H., Pinel, N., Arp, D. J., et al. (2010). *Nitrosopumilus maritimus* genome reveals unique mechanisms for nitrification and autotrophy in globally distributed marine crenarchaea. *Proc. Natl. Acad. Sci. U.S.A.* 107, 8818–8823. doi: 10.1073/pnas.0913531107
- Wallner, G., Amann, R., and Beisker, W. (1993). Optimizing fluorescent in situ hybridization with rRNA-targeted oligonucleotide probes for flow cytometric identification of microorganisms. *Cytometry* 14, 136–143. doi: 10.1002/cyto.990140205
- Wang, S., Xiao, X., Jiang, L., Peng, X., Zhou, H., Meng, J., et al. (2009). Diversity and abundance of ammonia-oxidizing archaea in hydrothermal vent chimneys of the Juan de Fuca Ridge. *Appl. Environ. Microbiol.* 75, 4216–4220. doi: 10.1128/AEM.01761-08
- Weidler, G. W., Dornmayr-Pfaffenhuemer, M., Gerbl, F. W., Heinen, W., and Stan-Lotter, H. (2007). Communities of archaea and bacteria in a subsurface radioactive thermal spring in the Austrian central Alps, and evidence of ammonia-oxidizing *Crenarchaeota*. *Appl. Environ. Microbiol.* 73, 259–270. doi: 10.1128/AEM.01570-06
- Wick, R. R., Judd, L. M., Gorrie, C. L., and Holt, K. E. (2017). Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLOS Comput. Biol.* 13:e1005595. doi: 10.1371/journal.pcbi.1005595
- Yan, J., Beattie, T. R., Rojas, A. L., Schermerhorn, K., Gristwood, T., Trinidad, J. C., et al. (2017). Identification and characterization of a heterotrimeric archaeal DNA polymerase holoenzyme. *Nat. Commun.* 8:15075. doi: 10.1038/ncomms15075
- Zeglin, L. H., Taylor, A. E., Myrold, D. D., and Bottomley, P. J. (2011). Bacterial and archaeal *amoA* gene distribution covaries with soil nitrification properties across a range of land uses. *Environ. Microbiol. Rep.* 3, 717–726. doi: 10.1111/j.1758-2229.2011.00290.x
- Zhang, C. L., Ye, Q., Huang, Z., Li, W., Chen, J., Song, Z., et al. (2008). Global occurrence of archaeal *amoA* genes in terrestrial hot springs. *Appl. Environ. Microbiol.* 74, 6417–6426. doi: 10.1128/AEM.00843-08
- Zhang, Y., Tian, Z., Liu, M., Shi, Z. J., Hale, L., Zhou, J., et al. (2015). High concentrations of the antibiotic spiramycin in wastewater lead to high abundance of ammonia-oxidizing archaea in nitrifying populations. *Environ. Sci. Technol.* 49, 9124–9132. doi: 10.1021/acs.est.5b01293
- Zhao, W., Song, Z., Jiang, H., Li, W., Mou, X., Romanek, C. S., et al. (2011). Ammonia-oxidizing archaea in Kamchatka hot springs. *Geomicrobiol. J.* 28, 149–159. doi: 10.1080/01490451003753076
- Zhou, J., Bruns, M. A., and Tiedje, J. M. (1996). DNA recovery from soils of diverse composition. *Appl. Environ. Microbiol.* 62, 316–322.
- Zorraquino, M. A., Althaus, R. L., Roca, M., and Molina, M. P. (2011). Heat treatment effects on the antimicrobial activity of macrolide and lincosamide antibiotics in milk. *J. Food Prot.* 74, 311–315. doi: 10.4315/0362-028X.JFP-10-297

**Conflict of Interest Statement:** MA and RK own and run DNASense, the sequencing center at which the metagenomes were sequenced and the bins assembled.

The other authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Daebeler, Herbold, Vierheilig, Sedlacek, Pjevac, Albertsen, Kirkegaard, de la Torre, Daims and Wagner. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.